

Leading neutrons from polarized pp collisions

B.Z. Kopeliovich^{*,†}, I.K. Potashnikova^{*}, Ivan Schmidt^{*} and J. Soffer^{**}

^{*}*Departamento de Física y Centro de Estudios Subatómicos,*

Universidad Técnica Federico Santa María, Casilla 110-V, Valparaíso, Chile

[†]*Institut für Theoretische Physik der Universität, Philosophenweg 19, 69120 Heidelberg, Germany*

^{**}*Department of Physics, Temple University, Philadelphia, PA 19122-6082, USA*

Abstract. We calculate the cross section and single-spin azimuthal asymmetry, $A_n(t)$ for inclusive neutron production in pp collisions at forward rapidities relative to the polarized proton. Absorptive corrections to the pion pole generate a relative phase between the spin-flip and non-flip amplitudes, which leads to an appreciable spin asymmetry. However, the asymmetry observed recently in the PHENIX experiment at RHIC at very small $|t| \sim 0.01 \text{ GeV}^2$ cannot be explained by this mechanism.

Keywords: neutrons, pions, polarization, absorption

PACS: 13.85.Ni, 11.80.Gw, 12.40.Nn, 11.80.Cr

INTRODUCTION

We are considering the reaction of inclusive neutron production $pp \rightarrow Xn$ at large Feynman x_F as was measured recently by the PHENIX experiment at RHIC [1] at $\sqrt{s} = 200 \text{ GeV}$. The preliminary data [2] are shown in Fig. 1. The measurements were




FIGURE 1. Single-spin asymmetry A_n in reaction $pp \rightarrow nX$ measured at $\sqrt{s} = 200 \text{ GeV}$ [2]

done with a transversely polarized proton beam and the neutron was detected in the very near forward (or backward) direction relative to the polarized beam with a transverse momentum p_T of the order of 100 MeV. An appreciable single spin asymmetry was found for forward neutrons, for events with neutron momenta larger than 30 GeV $A_N = (-9.0 \pm 0.6 \pm 0.9)\%$ with an additional scale uncertainty $\left(1.0 \begin{smallmatrix} +0.47 \\ -0.24 \end{smallmatrix}\right)$. Data also show

no dramatic variation with x_F . According to the Feynman scaling, one should not expect any strong energy dependence of $A_N(z)$.

At the same time, neutrons produced with $x_F < 0$ demonstrate small asymmetry, consistent with zero. This fact is explained by the so called Abarbanel-Gross theorem [3] which predicts zero asymmetry for particles produced in the fragmentation region of an unpolarized beam. This theorem was proven within the approximation of triple-Regge poles. Regge cuts breakdown this statement, but the corrections calculated in [4] turned out to be rather small, less than 1%.

The cross section of neutron production in pp collisions was calculated recently in [5] where it was found that previous calculations [6, 7], although agreed with the ISR data, grossly underestimated the magnitude of absorptive corrections. New measurements done recently show that the normalization of the ISR data is overestimated by factor two. This was observed in pp collisions in the NA49 experiment [8], and in ep [9] collisions. Here we concentrate on the spin dependence of neutron production in the fragmentation region of the polarized beam caused by the absorptive corrections.

ABSORPTIVE CORRECTIONS TO THE PION POLE

Pion exchange contribution to the amplitude of neutron production, $pp \rightarrow nX$ in Born approximation has the form

$$A_{p \rightarrow n}^B(\vec{q}, z) = \frac{1}{\sqrt{z}} \bar{\xi}_n [\sigma_3 \tilde{q}_L + \vec{\sigma} \cdot \vec{q}_T] \xi_p \phi^B(q_T, z), \quad (1)$$

where $\vec{\sigma}$ are Pauli matrices; $\xi_{p,n}$ are the proton or neutron spinors; $\vec{q}_T$ is the transverse component of the momentum transfer; $\tilde{q}_L = (1 - z)m_N$. Fractional momentum z is related to the invariant mass M_X of X , $z \approx 1 - M_X^2/s$ for $1 - z \ll 1$, where $\sqrt{s}$ is the c.m. energy of pp collision. The 4-momentum transfer squared t has the form, $t = -(\tilde{q}_L^2 + q_T^2)/z$.

The differential cross section of inclusive neutron production reads [11, 12, 5],

$$\begin{aligned} z \frac{d\sigma_{p \rightarrow n}^B}{dz dq_T^2} &= \frac{1}{s} |A_{p \rightarrow n}^B(\vec{q}_T, z)|^2 = \\ &= \left(\frac{\alpha'_\pi}{8} \right)^2 |t| G_{\pi^+ pn}^2(t) |\eta_\pi(t)|^2 (1 - z)^{1-2\alpha_\pi(t)} \sigma_{tot}^{\pi^+ p}(s' = M_X^2), \end{aligned} \quad (2)$$

where $\eta_\pi(t)$ is the signature factor, and $G_{\pi^+ pn}^2(t) = \exp(tR_1^2)$ is the pionic formfactor (see details in [5]). The results of calculation with the Born approximation, Eq. (2), at $\sqrt{s} = 200, 62.7$ and 30.6 GeV are depicted versus ISR data [13] in Fig. 2. The data are shown at two energies $\sqrt{s} = 30.6$ GeV and 62.7 GeV. Correspondingly, we used these energies in our calculations. One can see that the Born approximation considerably exceeds the data and leads to a wrong energy dependence.

Absorptive corrections, or initial/final state interactions look simpler and factorize in impact parameters. Therefore, we switch to a Fourier transformed amplitude (1) which




FIGURE 2. *Left panel:* Born approximation (dashed curve) for neutron production and ISR data [13] at $\sqrt{s} = 60.7 \text{ GeV}$ and $p_T = 0$. Two solid curves upper and bottom show the effect of absorptive corrections calculated in the dipole approach ($\times S^{(5q)}$) and in hadronic representation ($\times S^{(hadr)}$) respectively. *Right panel:* Energy dependence of inclusive neutron production. Three upper curves present the forward cross section at $\sqrt{s} = 30.6 \text{ GeV}$ (solid), 62.7 GeV (dashed) and 200 GeV (dotted-dashed) calculated in the Born approximation. These curves corrected for absorption ($\times S^{(5q)}$) are presented by three curves at the bottom. Data at $\sqrt{s} = 30.6 \text{ GeV}$ and 62.7 GeV [13] are depicted by squares and triangles respectively

has the form,

$$f_{p \rightarrow n}^B(\vec{b}, z) = \frac{1}{\sqrt{z}} \bar{\xi}_n \left[\sigma_3 \tilde{q}_L \theta_0^B(b, z) - i \frac{\vec{\sigma} \cdot \vec{b}}{b} \theta_s^B(b, z) \right] \xi_p. \quad (3)$$

Here

$$\begin{aligned} \theta_0^B(b, z) &= \int d^2q e^{i\vec{b}\vec{q}} \phi^B(q_T, z) \\ &= N(z) \left\{ i \frac{\pi \alpha'_\pi}{2z\beta^2} K_0(b/\beta) + \frac{1}{1 - \beta^2 \epsilon^2} [K_0(\epsilon b) - K_0(b/\beta)] \right\}; \end{aligned} \quad (4)$$

$$\begin{aligned} \theta_s^B(b, z) &= \frac{1}{b} \int d^2q e^{i\vec{b}\vec{q}} (\vec{b} \cdot \vec{q}) \phi^B(q_T, z) \\ &= N(z) \left\{ i \frac{\pi \alpha'_\pi}{2z\beta^3} K_1(b/\beta) + \frac{1}{1 - \beta^2 \epsilon^2} \left[\epsilon K_1(\epsilon b) - \frac{1}{\beta} K_1(b/\beta) \right] \right\}; \end{aligned} \quad (5)$$

and

$$\begin{aligned} N(z) &= \frac{1}{2} g_{\pi^+pn} z(1-z) \alpha'_\pi (m_\pi^2 + q_L^2/z) e^{-R_1^2 q_L^2/z} A_{\pi p \rightarrow X}(M_X^2) \\ \epsilon^2 &= q_L^2 + z m_\pi^2, \\ \beta^2 &= \frac{1}{z} [R_1^2 - \alpha'_\pi \ln(1-z)]. \end{aligned} \quad (6)$$

The process under consideration at large $z \rightarrow 1$ is associated with creation of a large rapidity gap (LRG), $\Delta y = |\ln(1-z)|$, where no particle is produced. Absorptive corrections may be also interpreted as a suppression related to the survival probability of LRG.

To identify the projectile Fock states responsible for absorptive corrections we start with Fig. 3a containing the amplitude of pion-proton inelastic collision $\pi + p \rightarrow X$. The latter is usually described as color exchange leading to creation of two color octet states with a large rapidity interval $\sim \ln(M_X^2/s_0)$ ($s_0 = 1 \text{ GeV}^2$) as is illustrated in Fig. 3b. Perturbatively, the interaction is mediated by gluonic exchanges. Nonperturbatively, e.g.




FIGURE 3. **a:** Born graph with single pion exchange and excitation of the projectile proton, $p + \pi \rightarrow X$; **b:** inelastic proton-pion interaction interaction, $p + \pi \rightarrow X$, via color exchange leading to production of two color-octet dipoles which hadronize further to X ; **c:** Fock state representation of the previous mechanism. A color octet-octet dipole which is a 5-quark Fock component of the projectile proton, interacts with the target proton via π^+ exchange. This 5-quark state may experience initial and final state interaction via vacuum quantum number (Pomeron) exchange with the nucleons (ladder-like strips).

in the string model, hadron collision looks like crossing and flip of the strings.

According to Fig. 3b one may think that this is the produced color octet-octet state which experiences final state interactions with the recoil neutron. On the other hand, at high energies multiple interactions become coherent, and both initial and final state interactions must be included. This leads to a specific space-time development of the process at high energies, namely, the incoming proton fluctuates into a 5-quark state $|\{3q\}_8\{\bar{q}q\}_8\rangle$ long in advance of the interaction with the target via pion exchange, as is illustrated in Fig. 3c.

This picture can be also interpreted in terms of the Reggeon calculus [5], which helps to identify what was overlooked in the calculation of absorptive corrections done in [6].

The resulting amplitude exposes both initial and final state attenuation of the 5-quark state,

$$f_{p \rightarrow n}(b, z) = f_{p \rightarrow n}^B(b, z) S(b, z), \quad (7)$$

which are incorporated via the survival amplitude $S(b, z)$.

We evaluate the survival amplitude $S(b, z)$ within two models based on the color-dipole and hadronic representations.

Color-dipole model.

According to the dual parton model [14] one can present the survival amplitude of a 5-quark state with an accuracy $1/N_c^2$ as,

$$S^{(5q)}(b) = S^{(3q)}(b) S^{(q\bar{q})}(b) = \left[1 - \text{Im} \Gamma^{(3q)p}(b)\right] \left[1 - \text{Im} \Gamma^{(q\bar{q})p}(b)\right]. \quad (8)$$

similar to Eq. (11), The elastic amplitudes $\Gamma^{(\bar{3}3)p}(b)$ of a color $\{\bar{3}3\}$ dipole interacting with a proton can be calculated in terms of the partial dipole elastic amplitudes, for which a saturated model was proposed recently [15, 5]. The results for $S^{(3q)}(b)$, $S^{(q\bar{q})}(b)$ and their product $S^{(5q)}(b)$ at $\sqrt{s} = 60 \text{ GeV}$ and $z = 0.8$ are depicted in Fig. 4 (left panel).




FIGURE 4. Partial survival amplitude $S(b, z)$ at $\sqrt{s} = 60 \text{ GeV}$ and $z = 0.8$. *Left panel:* Survival amplitudes $S^{(2q)}(b, z)$ and $S^{(3q)}(b, z)$ depicted by dot-dashed and dashed curves, respectively. Solid curve corresponds to $S^{(5q)}(b, z)$. *Right panel:* Survival amplitudes evaluated in hadronic representation.

Hadronic representation.

One can expand the 5-quark Fock state over the hadronic basis,

$$|\{3q\}_8\{\bar{q}q\}_8\rangle = d_0|p\rangle + d_1|N\pi\rangle + d_2|N2\pi\rangle + \dots \quad (9)$$

Since the admixture of sea quarks in the proton is small, the projection of the 5-quark state to the proton must be small too. It is natural to assume that the amplitude d_1 is the dominant one, since both states $|\{3q\}_8\{\bar{q}q\}_8\rangle$ and $|N\pi\rangle$ have the same valence quark content. Correspondingly, the absorption corrected partial amplitude gets the form,

$$f_{p \rightarrow n}(b, z) = f_{p \rightarrow n}^B(b, z) S^{(hadr)}(b), \quad (10)$$

where

$$S^{(hadr)}(b) = S^{\pi p}(b) S^{pp}(b) = [1 - \text{Im} \Gamma^{pp}(b)] [1 - \text{Im} \Gamma^{\pi p}(b)]. \quad (11)$$

The elastic partial amplitudes $\text{Im} \Gamma^{hp}(b)$ were extracted in a model independent way directly from data on elastic scattering [16, 5].

Our results for $S^{(hadr)}(b, z)$ are depicted in Fig. 4, right panel. The survival amplitude suppression factor $S^{(hadr)}(B, z)$ is rather similar to one calculated in the dipole model, $S^{(51)}(b, z)$.

ABSORPTION CORRECTED SPIN AMPLITUDES

An example of absorption corrected impact-parameter-dependent spin amplitudes is shown in Fig. 5 for $\sqrt{s} = 60 \text{ GeV}$ and $z = 0.8$ in comparison with the Born approximation.




FIGURE 5. Partial spin amplitudes for neutron production. Solid curves show the result of Born approximation. Dashed and dot-dashed curves include absorptive corrections calculated in the dipole approach ($\times S^{(5q)}(b, z)$) and in hadronic model ($\times S^{(hadr)}(b, z)$), respectively.

Now, it is straightforward to Fourier transform the absorption corrected amplitudes (7) back to momentum representation,

$$A_{p \rightarrow n}(\vec{q}, z) = \frac{1}{\sqrt{z}} \bar{\xi}_n [\sigma_3 \tilde{q}_L \phi_0(q_T, z) + \vec{\sigma} \cdot \vec{q}_T \phi_s(q_T, z)] \xi_p, \quad (12)$$

where

$$\begin{aligned} \text{Re } \phi_0(q_T, z) &= \frac{N(z)}{2\pi(1-\beta^2\epsilon^2)} \int_0^\infty db b J_0(bq_T) \left[K_0(\epsilon b) - K_0\left(\frac{b}{\beta}\right) \right] S(b, z); \\ \text{Im } \phi_0(q_T, z) &= \frac{\alpha'_\pi N(z)}{4z\beta^2} \int_0^\infty db b J_0(bq_T) K_0\left(\frac{b}{\beta}\right) S(b, z); \end{aligned} \quad (13)$$

$$\begin{aligned} q_T \text{Re } \phi_s(q_T, z) &= \frac{N(z)}{2\pi(1-\beta^2\epsilon^2)} \int_0^\infty db b J_1(bq_T) \left[\epsilon K_1(\epsilon b) - \frac{1}{\beta} K_1\left(\frac{b}{\beta}\right) \right] S(b, z); \\ q_T \text{Im } \phi_s(q_T, z) &= \frac{\alpha'_\pi N(z)}{4z\beta^3} \int_0^\infty db b J_1(bq_T) K_1\left(\frac{b}{\beta}\right) S(b, z). \end{aligned} \quad (14)$$

The forward neutron production cross section corrected for absorption is compared with the ISR data [13] in Fig. 2. The two models for absorption, dipole and hadronic,

are presented on the left panel by solid curves, upper and bottom respectively. The results of both models are pretty close to each other, but substantially underestimate the ISR data. It was argued, however, in [5] that the normalization of these data is twice overestimated. This is supported by recent direct measurements of $pp \rightarrow nX$ by the NA49 experiment [8] and by comparison with neutron production in DIS [9] (see discussion in [5]). Otherwise, the shape of the z -dependence is reproduced pretty well. Also the energy dependence of the cross section is much improved after inclusion of absorption. Apparently, the steep rise of the cross section with energy observed in Born approximation is nearly compensated by the falling energy dependence of LRG survival amplitudes. The results for q_T dependent differential cross section can be found in [5].

As far as the spin amplitudes are known, one can calculate the single spin asymmetry,

$$A_N(q_T, z) = \frac{2q_T q_L \phi_0(q_T, z) \phi_s(q_T, z)}{q_L^2 |\phi_0(q_T, z)|^2 + q_T^2 |\phi_s(q_T, z)|^2} \sin(\delta_0 - \delta_s), \quad (15)$$

where

$$\tan \delta_{0,s} = \frac{\text{Im} \phi_{0,s}(q_T, z)}{\text{Re} \phi_{0,s}(q_T, z)}. \quad (16)$$

Our predictions for $A_N(q_T, z)$ are presented in Fig. 6. Notice that the spin asymmetry




FIGURE 6. The curves in the first and second left panels correspond to the absorption corrections calculated in the dipole and hadronic models respectively. The spin asymmetry is calculated at five different fixed angles between the neutron and initial proton momenta in the c.m. of collision $\theta_{c.m.} = 0.001 - 0.005$. Correspondingly, neutron transverse and longitudinal momenta correlate, $q_T = \theta_{c.m.} z \sqrt{s}/2$. The last panel shows the q_T dependence of A_N calculated in the dipole model at four fixed values of $z = 0.6 - 0.9$.

depicted in the two left panels, demonstrate practically no model dependence of the spin asymmetry.

Unfortunately, our calculations at transverse momenta as small as was measured at RHIC [1] considerably underestimate the data. This is not a surprise, $q_T^2 \sim 0.01 \text{ GeV}^2$ is so small, that hadronic spin effects in other reactions are usually vanishingly small. The spin asymmetry which arises due to pion pole with absorptive corrections, develops a double zero at $q_T \rightarrow 0$. This is because both, the spin-flip amplitude and the relative phase, Eq. (16), vanish. However, at larger transverse momenta the single spin asymmetry reaches an appreciable value. One should include other mechanisms, for instance

interference of pion and a_1 exchanges, which is enhanced by a proper phase shift, $\pi/2$ and energy independence of the diffractive amplitude $\pi p \rightarrow a_1 p$.

ACKNOWLEDGMENTS

This work was supported in part by Fondecyt (Chile) grants 1050519 and 1050589, and by DFG (Germany) grant PI182/3-1.

REFERENCES

1. A. Bazilevsky *et al.*, Phys. Lett. B **650**, 325 (2007)
2. M. Togawa for the PHENIX collaboration, talk at the Conference "Spin 2006", Kyoto, Japan, October 2-7, 2006.
3. H. D. I. Abarbanel and D. J. Gross, Phys. Rev. Lett. **26**, 732 (1971); Phys. Rev. D **5**, 699 (1972).
4. B. Z. Kopeliovich, L. I. Lapidus and A. B. Zamolodchikov, Sov. J. Nucl. Phys. **26**, 209 (1977) [Yad. Fiz. **26**, 399 (1977)].
5. B. Z. Kopeliovich, I. K. Potashnikova, I. Schmidt and J. Soffer, arXiv:0805.4534 [hep-ph], to appear in Phys. Rev. D.
6. N. N. Nikolaev, W. Schafer, A. Szczurek and J. Speth, Phys. Rev. D **60**, 014004 (1999).
7. U. D'Alesio and H. J. Pirner, Eur. Phys. J. A **7**, 109 (2000).
8. D. Varga, NA49 Collaboration, Eur. Phys. J. C **33**, S515 (2004).
9. S. Chekanov *et al.* [ZEUS Collaboration], Nucl. Phys. B **637**, 3 (2002).
10. B. Kopeliovich, B. Povh and I. Potashnikova, Z. Phys. C **73**, 125 (1996).
11. M. Bishari, Phys. Lett. B **38**, 510 (1972).
12. Yu. M. Kazarinov, B. Z. Kopeliovich, L. I. Lapidus and I. K. Potashnikova, Sov. Phys. JETP **43**, 598 (1976) [Zh. Eksp. Teor. Fiz. **70**, 1152 (1976)].
13. W. Flauger and F. Mönig, Nucl. Phys. B **109** (1976) 347.
14. A. Capella, U. Sukhatme, C. I. Tan and J. Tran Thanh Van, Phys. Rept. **236**, 225 (1994).
15. B. Z. Kopeliovich, H. J. Pirner, A. H. Rezaeian and I. Schmidt, Phys. Rev. D **77**, 034011 (2008).
16. B. Z. Kopeliovich, I. K. Potashnikova, B. Povh and E. Predazzi, Phys. Rev. Lett. **85**, 507 (2000); Phys. Rev. D **63**, 054001 (2001).